

## **NIDEM\_25\_1.0.0**

Submitted by robbibt on Wed, 03/10/2018 - 11:40

[Basic information](#)

Simple title

National Intertidal Digital Elevation Model

Descriptive title

National Intertidal Digital Elevation Model 25m 1.0.0

Version number

1.0.0

Abstract

The National Intertidal Digital Elevation Model (NIDEM) is a continental-scale elevation dataset for Australia's exposed intertidal zone. NIDEM provides the first three-dimensional representation of Australia's intertidal sandy beaches and shores, tidal flats and rocky shores and reefs at 25 m spatial resolution, addressing a key gap between the availability of sub-tidal bathymetry and terrestrial elevation data. NIDEM was generated by combining global tidal modelling with a 30-year time series archive of spatially and spectrally calibrated Landsat satellite data managed within the Digital Earth Australia (DEA) platform. NIDEM complements existing intertidal extent products, and provides data to support a new suite of use cases that require a more detailed understanding of the three-dimensional topography of the intertidal zone, such as hydrodynamic modelling, coastal risk management and ecological habitat mapping.

### General Public Description

The National Intertidal Digital Elevation Model (NIDEM) is the first 3D model of Australia's intertidal zone - the area of coastline exposed and flooded by ocean tides. Intertidal environments support important ecological habitats (e.g. tidal mudflats, sandy beaches and coral reefs), and provide many valuable benefits such as storm surge protection, carbon storage and natural resources for recreational and commercial use. However, intertidal zones are faced with increasing threats from coastal erosion, land reclamation (e.g. port construction), and sea level rise. Accurate elevation data describing the height and shape of the coastline is needed to help predict when and where these threats will have the greatest impact. However, this data is expensive and challenging to map across the entire intertidal zone of a continent the size of Australia.

The rise and fall of the ocean can be used to describe the three-dimensional shape of the coastline by mapping the location of the land-sea boundary (or 'waterline') across a range of known tides (e.g. low tide, high tide). To model the elevation of Australia's entire intertidal zone, we combined thousands of satellite images of the coastline taken at different tide height conditions between 1987 and 2016. Assuming that waterlines mapped from satellite data represent lines of constant height compared to mean sea level (MSL), these contours could be analysed to estimate the height of every 25 x 25 m area of coastline between the highest and lowest observed tide. The resulting NIDEM dataset provides a first-of-its kind source of elevation data for Australia's intertidal zone. It is expected that NIDEM will be used to map the habitats of threatened coastal species, identify areas of coastal erosion, plan for extreme events such as storm surges and flooding, and improve models of how sea level rise will affect the Australian coastline.

To view and access the data interactively, visit <http://nationalmap.gov.au>. Click 'Add data', and then the 'Surface Water and Marine > Coastal > Intertidal elevation model' theme. Click on 'NIDEM - Intertidal elevation model' and then 'Add to the map'.

Owner

[Commonwealth of Australia \(Geoscience Australia\)](#)

Authors/principal contributors

Robbi Bishop-Taylor

Stephen Sagar

Leo Lymburner

License

[CC BY Attribution 4.0 International License](#)

DOI

[eCat catalogue for data access](#)

[Scientific reference \(Bishop-Taylor et al. 2019\)](#)

Websites

[Digital Earth Australia website](#)

Services

[Published via WMS services available on this link with layer name: NIDEM](#)

Catalogues

[THREDDS](#)


[Geoscience Australia eCat](#)

Documents/References


[Internal Peer Review - David Gavin](#)

[Author Review - Robbi Bishop-Taylor](#)


Images





10% intervals of the tidal range mapped from 30 years of Landsat satellite imagery



Waterline contours extracted and assigned median tidal heights for each 10% interval




Interpolated elevation relative to mean sea level  
< 2.50 m      0.00 m      > 1.50 m



Elevation relative to modelled MSL





## Technical Details

### Overview

Intertidal environments support important ecological habitats (e.g. sandy beaches and shores, tidal flats and rocky shores and reefs), and provide many valuable benefits such as storm surge protection, carbon storage and natural resources for recreational and commercial use. However, intertidal zones are faced with increasing threats from coastal erosion, land reclamation (e.g. port construction), and sea level rise. Accurate elevation data describing the height and shape of the coastline is needed to help predict when and where these threats will have the greatest impact. However, this data is expensive and challenging to map across the entire intertidal zone of a continent the size of Australia.

The rise and fall of the ocean can be used to describe the three-dimensional shape of the coastline by mapping the land-sea boundary (or 'waterline') across a range of known tides (e.g. low tide, high tide). Assuming that these waterlines represent lines of constant height relative to mean sea level (MSL), elevations can be modelled for the area of coastline located between the lowest and highest observed tide. To model the elevation of Australia's entire intertidal zone, 30 years of satellite images of the coastline (between 1986 and 2016 inclusive) were obtained from the archive of spatially and spectrally calibrated Landsat observations managed within the Digital Earth Australia (DEA) platform. Using the improved tidal modelling framework of the Intertidal Extents Model v2.0 (ITEM 2.0; Sagar et al. 2017, 2018), each satellite observation in the 30 year time series could be accurately associated with a modelled tide height using the global TPX08 ocean tidal model. These satellite observations were converted into a water index (NDWI), composited into discrete ten percent intervals of the observed tide range (e.g. the lowest 10% of observed tides etc), and used to extract waterlines using a spatially consistent and automated waterline extraction procedure. Triangulated irregular network (TIN) interpolation was then used to derive elevations relative to modelled mean sea level for each 25 x 25 m Landsat pixel across approximately 15,387 sq. km of intertidal terrain along Australia's entire coastline.

NIDEM differs from previous methods used to model the elevation of the intertidal zone which have predominately focused on extracting waterlines from a limited selection of satellite images using manual digitisation and visual interpretation (e.g. Chen and Rau 1998; Zhao et al. 2008; Liu et al. 2013; Chen et al. 2016). This manual process introduces subjectivity, is impractical to apply at a continental-scale, and has inherent restrictions based on the availability of high quality image data at appropriate tidal stages. By developing an automated approach to generating satellite-derived elevation data based on a 30 year time series of observations managed within the Digital Earth Australia (DEA) platform, it was possible to produce the first continental-scale three-dimensional model of the intertidal zone.

### Features

The National Intertidal Digital Elevation Model (NIDEM) consists of four raster datasets (NIDEM, unfiltered NIDEM, NIDEM mask, NIDEM uncertainty) and two vector datasets (NIDEM waterline contours and ITEM v2.0 tidal modelling polygons). For convenience, the four raster datasets are provided both as individual geotiff/NetCDF files, and continental-scale geotiff mosaics.

## NIDEM

The NIDEM dataset provides elevation in metre units relative to modelled Mean Sea Level for each pixel of intertidal terrain across the Australian coastline. The DEMs have been cleaned by masking out non-intertidal pixels and pixels where tidal processes poorly explain patterns of inundation (see NIDEM

mask below). This is the primary output product, and is expected to be the default product for most applications. The dataset consists of 306 raster files corresponding to the 306 ITEM v2.0 tidal modelling polygons, and a continental-scale mosaic.

## **Unfiltered NIDEM**

The unfiltered NIDEM dataset provides un-cleaned elevation in metre units relative to modelled Mean Sea Level for each pixel of intertidal terrain across the Australian coastline. Compared to the default NIDEM product, these layers have not been filtered to remove noise, modelling artefacts or invalid elevation values (see NIDEM mask below). This supports applying custom filtering methods to the raw NIDEM data. The dataset consists of 306 raster files corresponding to the 306 ITEM v2.0 tidal modelling polygons, and a continental-scale mosaic.

## **NIDEM mask**

The NIDEM mask dataset flags non-intertidal terrestrial pixels with elevations greater than 25 m, and sub-tidal pixels with depths greater than -25 m relative to Mean Sea Level. Pixels where tidal processes poorly explain patterns of inundation are also flagged by identifying any pixels with ITEM confidence NDWI standard deviation greater than 0.25. The NIDEM mask was used to filter and clean the NIDEM dataset to remove modelling artefacts and noise (e.g. intertidal pixels in deep water or high elevations) and invalid elevation estimates caused by coastal change or poor model performance. The dataset consists of 306 raster files corresponding to the 306 ITEM v2.0 tidal modelling polygons, and a continental-scale mosaic.

## **NIDEM uncertainty**

The NIDEM uncertainty dataset provides a measure of the uncertainty (not to be confused with accuracy) of NIDEM elevations in metre units for each pixel. The range of Landsat observation tide heights used to compute median tide heights for each waterline contour can vary significantly between tidal modelling polygons. To quantify this range, the standard deviation of tide heights for all Landsat images used to produce each ITEM interval and subsequent waterline contour was calculated. These values were interpolated to return an estimate of uncertainty for each individual pixel in the NIDEM datasets: larger values indicate the waterline contour was based on a composite of images with a larger range of tide heights. The dataset consists of 306 raster files corresponding to the 306 ITEM v2.0 tidal modelling polygons, and a continental-scale mosaic.

## **NIDEM waterline contours**

The NIDEM waterline contour dataset provides a vector representation of the boundary of every ten percent interval of the observed intertidal range. These contours were extracted along the boundary between each ITEM v2.0 tidal interval, and assigned the median and standard deviation (see NIDEM uncertainty above) of tide heights from the ensemble of corresponding Landsat observations. These datasets facilitate re-analysis by allowing alternative interpolation methods (e.g. kriging, splines) to be used to generate DEMs from median tide heights. The dataset consists of 306 shapefiles corresponding to the 306 ITEM v2.0 tidal modelling polygons.

## **ITEM v2.0 tidal modelling polygons**

The ITEM v2.0 tidal model polygon dataset describes the multi-resolution tidal framework developed by Sagar et al. (2018). The framework uses partitioning methods to allow spatial variability in the tidal model to drive the size and locations of a Voronoi polygon mesh. The 306 resulting tidal modelling polygons are then used as analysis units for tidal modelling, with tide height predictions defined at the nodes of each Voronoi cell. To evaluate the representativeness of NIDEM data compared to the full tidal range, the spread of tidal heights coincident with the input Landsat imagery was compared against the full range of modelled tide heights present within each tidal modelling polygon. Three indices were calculated: spread (the proportion of the full modelled tidal range observed by Landsat), low tide offset (the proportion of the lowest tidal heights not observed by Landsat) and high tide offset (the proportion of the highest tidal heights not observed by Landsat). This evaluation of the observed tidal range at a particular location provides valuable information to users about the "fitness for purpose" of NIDEM at a given location for their specific application (see Bishop-Taylor et al. 2019).

### **Background (Lineage Statement)**

The product differs from previous methods used to model the elevation of the intertidal zone which have

predominately focused on extracting waterlines from a limited selection of satellite images using manual digitisation and visual interpretation (e.g. Chen and Rau 1998; Zhao et al. 2008; Liu et al. 2013; Chen et al. 2016). This manual process introduces subjectivity, is impractical to apply at a continental-scale, and has inherent restrictions based on the availability of high quality image data at appropriate tidal stages. By developing an automated approach to generating satellite-derived elevation data based on a 30 year time series of observations managed within the Digital Earth Australia (DEA) platform, we were able to produce the first continental-scale three-dimensional model of the intertidal zone.

## Applications

By providing an important 'missing link' between existing terrestrial topographic and marine bathymetric datasets, we anticipate elevation data from NIDEM will contribute to unified coastal terrain models that combine best-available topography and bathymetry datasets into single continuous 'topobathy' datasets. This would provide coastal managers with near-seamless elevation data extending from inland to the ocean floor, and support a holistic approach to understanding the physical and ecological processes influencing the coastal zone. This may include providing valuable baseline elevation data for predicting the impact of coastal hazards such as storm surges or tsunami inundation, investigating mechanisms of coastal erosion and sediment transport, or improving modelling of sea-level rise under future climate change scenarios.

Access to accurate intertidal elevation data is also critical for studying and conserving coastal ecosystems, particularly given the important role intertidal topography and tidal dynamics play in regulating environmental stress and spatial patterns of species richness, abundance and productivity. Data from NIDEM will facilitate whole-of-landscape approaches to ecological modelling and three-dimensional habitat mapping in the intertidal zone that are not artificially restricted to either marine or terrestrial ecosystems.

## Tags

[EGD](#)  
[NEMO](#)  
[DEA](#)

## Accuracy and limitations

### Accuracy

To assess the accuracy of NIDEM, we compared modelled elevations against three independent elevation and bathymetry validation datasets: the DEM of Australia derived from LiDAR 5 Metre Grid (Geoscience Australia, 2015), elevation data collected from Real Time Kinematic (RTK) GPS surveys (Danaher & Collett, 2006; HydroSurvey Australia, 2009), and 1.0 m resolution multibeam bathymetry surveys (Solihuddin et al., 2016). We assessed overall accuracy across three distinct intertidal environments: sandy beaches and shores, tidal flats, and rocky shores and reefs:

- Sandy beaches and shores, 5 sites: Pearson's correlation = 0.92, Spearman's correlation = 0.93, RMSE +/- 0.41 m
- Tidal flats, 9 sites: Pearson's correlation = 0.78, Spearman's correlation = 0.81, RMSE +/- 0.39 m
- Rocky shores and reefs, 7 sites: Pearson's correlation = 0.46, Spearman's correlation = 0.79, RMSE +/- 2.98 m

### Limitations

NIDEM covers the exposed intertidal zone which includes sandy beaches and shores, tidal flats and rocky shores and reefs. The model excludes intertidal vegetation communities such as mangroves.

Areas with comparatively steep coastlines and small tidal ranges are poorly captured in the 25 m spatial resolution input Landsat data and resulting NIDEM model. This includes much of the south eastern and southern Australian coast (e.g. New South Wales, Victoria, Tasmania).

Poor validation results for rocky shore and reef sites within the southern Kimberly region highlighted limitations in the NIDEM model that occur when the global OTPS TPX08 Atlas Tidal Model was unable to predict complex and asynchronous local tidal patterns. This is likely to also reduce model accuracy in complex estuaries and coastal wetlands where river flow or vegetative resistance causes hydrodynamic attenuation in tidal flow.

The complex temporal behaviour of tides mean that a sun synchronous sensor like Landsat does not observe the full range of the tidal cycle at all locations. This causes spatial bias in the proportion of the



tidal range observed in different regions, which can prevent NIDEM from providing elevation data for areas of the intertidal zone exposed or inundated at the extremes of the tidal range. Accordingly, NIDEM provides elevation data for the portion of the tidal range observed by Landsat, rather than the full tidal range.

While image compositing and masking methods have been applied to remove the majority of noise and non-tidal artefacts from NIDEM, issues remain in several locations. It is recommended that the data be used with caution in the following areas:

- The Recherche Archipelago in southern Western Australia
- Port Phillip Bay in Victoria
- The eastern coast of Tasmania and King Island
- Saunders Reef and surrounds in the northern Coral Sea

Update frequency

[As needed](#)

Data sources

[ITEM\\_25\\_2.0.0](#)

[High-resolution depth model for Northern Australia - 30 m](#)

[SR-NT\\_25\\_2.0.0](#)

Major algorithms

[OTPS TPX08 Atlas Tidal Model](#)

[Waterline contour extraction](#)

[Triangulated Irregular Network interpolation](#)

Processing steps

[Extract waterline contours along boundary of each ITEM tidal interval \(i.e. each ten percent of the observed tidal range\)](#)

[Assign contours with median and standard deviation tide heights of Landsat observations for each tidal interval](#)

[Generate continuous elevation data from median tide heights per contour using Triangulated Irregular Network interpolation](#)

[Clean and filter DEMs to remove noise, modelling artefacts and invalid elevation estimates](#)

Custodian

[Commonwealth of Australia \(Geoscience Australia\)](#)

Acknowledgments

We thank the individuals and organisations that provided validation data for this study. RTK GPS data used in Darwin Harbour, NT was provided by HydroSurvey Australia, with the permission of GHD and Power & Water Corporation, NT. RTK GPS data used in Moreton Bay, QLD was provided by Tim Danaher, NSW Office of Environment and Heritage. Multibeam bathymetry data over the Kimberley reefs was provided by M. O'Leary at Curtin University, WA. We thank C. Burton, S. Chua, A. Haiblen, A. Lem, B. Moushall, B. Stepin and S. Wong for assisting with field validation. This research was undertaken with the assistance of resources from the National Computational Infrastructure High Performance Data (HPD) platform, which is supported by the Australian Government.

Subject matter experts

[Robbi Bishop-Taylor](#)

[Stephen Sagar](#)

Rights statement

© Commonwealth of Australia (Geoscience Australia) 2015. Creative Commons Attribution 4.0 International License. <https://creativecommons.org/licenses/by/4.0/>.

Quality assurance

To assess the accuracy of NIDEM, we compared modelled elevations against three independent elevation and bathymetry validation datasets: the DEM of Australia derived from LiDAR 5 Metre Grid (Geoscience Australia, 2015), elevation data collected from Real Time Kinematic (RTK) GPS surveys (Danaher & Collett, 2006; HydroSurvey Australia, 2009), and 1.0 m resolution multibeam bathymetry surveys (Solihuddin et al., 2016). We assessed overall accuracy across three distinct intertidal environments: sandy beaches and shores, tidal flats, and rocky shores and reefs:

- Sandy beaches and shores, 5 sites: Pearson's correlation = 0.92, Spearman's correlation = 0.93, RMSE +/- 0.41 m

- Tidal flats, 9 sites: Pearson's correlation = 0.78, Spearman's correlation = 0.81, RMSE +/- 0.39 m
- Rocky shores and reefs, 7 sites: Pearson's correlation = 0.46, Spearman's correlation = 0.79, RMSE +/- 2.98 m

## Data description

The National Intertidal Digital Elevation Model (NIDEM) consists of four raster datasets (NIDEM, unfiltered NIDEM, NIDEM mask, NIDEM uncertainty) and two vector datasets (NIDEM waterline contours and ITEM v2.0 tidal modelling polygons). For convenience, the four raster datasets are provided both as individual geotiff/NetCDF files, and continental-scale geotiff mosaics.

## NIDEM

The NIDEM dataset provides elevation in metre units relative to modelled Mean Sea Level for each pixel of intertidal terrain across the Australian coastline. The DEMs have been cleaned by masking out non-intertidal pixels and pixels where tidal processes poorly explain patterns of inundation (see NIDEM mask below). This is the primary output product, and is expected to be the default product for most applications. The dataset consists of 306 raster files corresponding to the 306 ITEM v2.0 tidal modelling polygons, and a continental-scale mosaic.

### *Naming convention:*

- NIDEM\_{polygon\_id}\_{tidalmodelling\_coords}
- (e.g. NIDEM\_33\_130.91\_-12.26.tif)
- Continental mosaic: NIDEM\_mosaic.tif

### *Attributes:*

- Single band float raster
- -9999: NoData
- All other values: elevation in metre units relative to modelled Mean Sea Level

## Unfiltered NIDEM

The unfiltered NIDEM dataset provides un-cleaned elevation in metre units relative to modelled Mean Sea Level for each pixel of intertidal terrain across the Australian coastline. Compared to the default NIDEM product, these layers have not been filtered to remove noise, modelling artefacts or invalid elevation values (see NIDEM mask below). This supports applying custom filtering methods to the raw NIDEM data. The dataset consists of 306 raster files corresponding to the 306 ITEM v2.0 tidal modelling polygons, and a continental-scale mosaic.

### *Naming convention:*

- NIDEM\_unfiltered\_{polygon\_id}\_{tidalmodelling\_coords}
- (e.g. NIDEM\_unfiltered\_33\_130.91\_-12.26.tif)
- Continental mosaic: NIDEM\_unfiltered\_mosaic.tif

### *Attributes:*

- Single band float raster
- -9999: NoData
- All other values: elevation in metre units relative to modelled Mean Sea Level

## NIDEM mask

The NIDEM mask dataset flags non-intertidal terrestrial pixels with elevations greater than 25 m, and sub-tidal pixels with depths greater than -25 m relative to Mean Sea Level. Pixels where tidal processes poorly explain patterns of inundation are also flagged by identifying any pixels with ITEM confidence NDWI standard deviation greater than 0.25. The NIDEM mask was used to filter and clean the NIDEM dataset to remove modelling artefacts and noise (e.g. intertidal pixels in deep water or high elevations) and invalid elevation estimates caused by coastal change or poor model performance. The dataset consists of 306 raster files corresponding to the 306 ITEM v2.0 tidal modelling polygons, and a

continental-scale mosaic.

*Naming convention:*

- NIDEM\_mask\_{polygon\_id}\_{tidalmodelling\_coords}
- (e.g. NIDEM\_mask\_33\_130.91\_-12.26.tif)
- Continental mosaic: NIDEM\_mask\_mosaic.tif

*Attributes:*

- Single band integer raster
- 1: Non-intertidal terrestrial pixels with elevations greater than 25 m
- 2: Sub-tidal pixels with depths greater than -25 m
- 3: ITEM confidence NDWI standard deviation greater than 0.25 (pixels where tidal processes poorly explain patterns of inundation)
- -9999: NoData

## **NIDEM uncertainty**

The NIDEM uncertainty dataset provides a measure of the uncertainty (not to be confused with accuracy) of NIDEM elevations in metre units for each pixel. The range of Landsat observation tide heights used to compute median tide heights for each waterline contour can vary significantly between tidal modelling polygons. To quantify this range, the standard deviation of tide heights for all Landsat images used to produce each ITEM interval and subsequent waterline contour was calculated. These values were interpolated to return an estimate of uncertainty for each individual pixel in the NIDEM datasets: larger values indicate the waterline contour was based on a composite of images with a larger range of tide heights. The dataset consists of 306 raster files corresponding to the 306 ITEM v2.0 tidal modelling polygons, and a continental-scale mosaic.

*Naming convention:*

- NIDEM\_uncertainty\_{polygon\_id}\_{tidalmodelling\_coords}
- (e.g. NIDEM\_uncertainty\_33\_130.91\_-12.26.tif)
- Continental mosaic: NIDEM\_uncertainty\_mosaic.tif

*Attributes:*

- Single band float raster
- -9999: NoData
- All other values: standard deviation of tide heights of all Landsat imagery used to generate each ITEM tidal interval in metre units

## **NIDEM waterline contours**

The NIDEM waterline contour dataset provides a vector representation of the boundary of every ten percent interval of the observed intertidal range. These contours were extracted along the boundary between each ITEM v2.0 tidal interval, and assigned the median and standard deviation (see NIDEM uncertainty above) of tide heights from the ensemble of corresponding Landsat observations. These datasets facilitate re-analysis by allowing alternative interpolation methods (e.g. kriging, splines) to be used to generate DEMs from median tide heights. The dataset consists of 306 shapefiles corresponding to the 306 ITEM v2.0 tidal modelling polygons.

*Naming convention:*

- NIDEM\_contours\_{polygon\_id}\_{tidalmodelling\_coords}
- (e.g. NIDEM\_contours\_33\_130.91\_-12.26.shp)

*Attributes:*

- Polyline shapefile
- elev\_m: median tide height of the ensemble of Landsat observations for each ten percent tidal interval



- uncert\_m: standard deviation tide height of the ensemble of Landsat observations for each ten percent tidal interval

## ITEM v2.0 tidal modelling polygons

The ITEM v2.0 tidal model polygon dataset describes the multi-resolution tidal framework developed by Sagar et al. (2018). The framework uses partitioning methods to allow spatial variability in the tidal model to drive the size and locations of a Voronoi polygon mesh. The 306 resulting tidal modelling polygons are then used as analysis units for tidal modelling, with tide height predictions defined at the nodes of each Voronoi cell. To evaluate the representativeness of NIDEM data compared to the full tidal range, the spread of tidal heights coincident with the input Landsat imagery was compared against the full range of modelled tide heights present within each tidal modelling polygon. Three indices were calculated: spread (the proportion of the full modelled tidal range observed by Landsat), low tide offset (the proportion of the lowest tidal heights not observed by Landsat) and high tide offset (the proportion of the highest tidal heights not observed by Landsat). This evaluation of the observed tidal range at a particular location provides valuable information to users about the "fitness for purpose" of NIDEM at a given location for their specific application (see Bishop-Taylor et al. 2019).

*Naming convention:*

- ITEMv2\_tidalmodel.shp

*Attributes:*

- Polygon shapefile
- polygon\_id: tidal modelling polygon ID (ranges from 1 to 306)
- lat: latitude of the Voronoi point used to derive tidal heights for polygon
- lon: longitude of the Voronoi point used to derive tidal heights for polygon
- spread: proportion of the full modelled tidal range observed by Landsat for polygon
- offset\_low: low tide offset; the proportion of the lowest tidal heights not observed by Landsat for polygon
- offset\_high: high tide offset; the proportion of the highest tidal heights not observed by Landsat for polygon

Schema/Spatial Extent

## DEA Albers Raster Schema

Update frequency

[As needed](#)

Temporal extent

Sat, 16/08/1986 - 00:00 - Mon, 31/07/2017 - 23:59

Spatial extent

Minimum X value

-1943830.00

Maximum X value

2170690.00

Minimum Y value

-1119030.00

Maximum Y value

-4856630.00

Cell size X

25.00

Cell size Y

25.00

Pixel origin

[Top Left](#)

Coordinate Reference System

[Australian Albers / GDA94 \(EPSG: 3577\)](#)

Coordinate System Units

[metre](#)

[Management/Reporting](#)

## Intended Delivery Channels

Data staging location

[http://dapds00.nci.org.au/thredds/catalogs/fk4/nidem\\_1\\_0.html](http://dapds00.nci.org.au/thredds/catalogs/fk4/nidem_1_0.html)

Data final location/folder name

[http://dapds00.nci.org.au/thredds/catalogs/fk4/nidem\\_1\\_0.html](http://dapds00.nci.org.au/thredds/catalogs/fk4/nidem_1_0.html)

Web Service Data Description/Abstract

The National Intertidal Digital Elevation Model (NIDEM) product is a continental-scale dataset providing continuous elevation data for Australia's exposed intertidal zone. NIDEM provides the first three-dimensional representation of Australia's intertidal zone (excluding off-shore Territories and intertidal mangroves) at 25 m spatial resolution, addressing a key gap between the availability of sub-tidal bathymetry and terrestrial elevation data. NIDEM was generated by combining global tidal modelling with a 30-year time series archive of spatially and spectrally calibrated Landsat satellite data managed within the Digital Earth Australia (DEA) platform. NIDEM complements existing intertidal extent products, and provides data to support a new suite of use cases that require a more detailed understanding of the three-dimensional topography of the intertidal zone, such as hydrodynamic modelling, coastal risk management and ecological habitat mapping. For service status information, see <https://status.dea.ga.gov.au>

One line summary

The National Intertidal Digital Elevation Model (NIDEM) product is a continental-scale elevation dataset for Australia's exposed intertidal zone

Security classification

## Unclassified

eCat ID

123678

eCat URL

<http://pid.geoscience.gov.au/dataset/ga/123678>

HPRM PMP references

D2018-158245

HPRM Approval Reference

D2018-158044

Projects

## National Intertidal Digital Elevation Model

### Contacts

Principal contact

[Geoscience Australia - Client Services](#)

Publishing entity

[Commonwealth of Australia \(Geoscience Australia\)](#)

Ordering entity

[Geoscience Australia - Client Services](#)

Distribution entity

[Commonwealth of Australia \(Geoscience Australia\)](#)

Ingestion contact

[Geoscience Australia - Team Leader, Applications and Systems Support, NEMO](#)

QA contact

[Geoscience Australia - Team Leader, Business Systems Development Section, NEMO](#)

Data source contact

[Geoscience Australia - Team Leader, Applications and Systems Support, NEMO](#)

Maintenance contact

[Geoscience Australia - Team Leader, Applications and Systems Support, NEMO](#)

Metadata contact

[Geoscience Australia - Team Leader, Product Management, Operations Section NEMO](#)

Media contact

[Geoscience Australia - GA Media Hotline](#)

Feedback contact

[Geoscience Australia - Section Leader, Operations Section, NEMO](#)

THREDDS

On